

# A Characterization of Banach Spaces Containing $l^1$

(pointwise convergence/Boolean independence)

HASKELL P. ROSENTHAL

The Ohio State University, Columbus, Ohio 43210; and The University of California, Berkeley, Calif. 94720

Communicated by Charles B. Morrey, Jr., March 29, 1974

**ABSTRACT** It is proved that a Banach space contains a subspace isomorphic to  $l^1$  if (and only if) it has a bounded sequence with no weak-Cauchy subsequence. The proof yields that a sequence of subsets of a given set has a subsequence that is either convergent or Boolean independent.

A bounded sequence of elements  $(f_n)$  in a Banach space  $B$  is said to be *equivalent to the usual  $l^1$ -basis* provided there is a  $\delta > 0$  so that for all  $n$  and choices of scalars  $c_1, \dots, c_n$ ,

$$\delta \sum_{i=1}^n |c_i| \leq \|\sum_{i=1}^n c_i f_i\|. \quad [1]$$

Of course if  $(f_n)$  has this property, then the closed linear span of the  $f_n$ 's is isomorphic (linearly homeomorphic) to  $l^1$ .  $(f_n)$  is said to be a *weak-Cauchy sequence* if  $\lim_{n \rightarrow \infty} b^*(f_n)$  exists for all

$b^* \in B^*$ , the dual of  $B$ .

**THE MAIN THEOREM.** Let  $(f_n)$  be a bounded sequence in a real Banach space  $B$ . Then  $(f_n)$  has a subsequence  $(f'_n)$  satisfying one of the following two mutually exclusive alternatives:

- (i)  $(f'_n)$  is a weak-Cauchy sequence.
- (ii)  $(f'_n)$  is equivalent to the usual  $l^1$ -basis.

We note two immediate consequences:

I. If  $B$  is weakly complete (that is, every weak-Cauchy sequence in  $B$  converges weakly to an element of  $B$ ), then  $B$  is either reflexive or contains a subspace isomorphic to  $l^1$ .

II. If  $B$  has the Schur property (that is, every weakly convergent sequence in  $B$  converges in norm), then every infinite-dimensional subspace of  $B$  contains a subspace isomorphic to  $l^1$ .

It is a well-known consequence of the Vitali-Hahn-Saks theorem that  $L^1(\mu)$  is weakly complete for any measure  $\mu$  on a measurable space, while  $l^1$  has the Schur property.

We reformulate the *Main Theorem* as follows:

**THEOREM 1.** Let  $S$  be a set and  $(f_n)$  a uniformly bounded sequence of real-valued functions defined on  $S$ . Then  $(f_n)$  has a subsequence  $(f'_n)$  satisfying one of the following alternatives:

- (i)  $(f'_n)$  converges point-wise on  $S$ .
- (ii)  $(f'_n)$  is equivalent in the supremum norm to the usual  $l^1$ -basis.

The exclusivity of the alternatives of the *Main Theorem* follows from the obvious fact that the usual  $l^1$ -basis is not a weak-Cauchy sequence. If  $(b_n)$  is a bounded sequence in a Banach space  $B$ , we let  $S$  denote the unit ball of  $B^*$  and then define  $f_n(s) = s(b_n)$  for all  $s \in S$  and  $n$ , to deduce the *Main Theorem* from *Theorem 1*.

We begin the proof of *Theorem 1* with that of the crucial special case of characteristic functions; that is, a sequence  $(A_n)$  of subsets of  $S$  with  $f_n = \chi_{A_n}$  for all  $n$  (where  $\chi_{A_n}(s) = 1$  if  $s \in A_n$ ;  $\chi_{A_n}(s) = 0$  if  $s \notin A_n$ ). (In classical terminology,  $(A_n)$  is said to converge if  $\chi_{A_n}$  converges point-wise.) Our proof of this special case yields that if  $(A_n)$  has no convergent subsequence, then  $(A_n)$  has a Boolean independent subsequence

$(A_{n'})$ ; that is, for every pair of nonempty finite disjoint subsets  $G$  and  $B$  of indices,  $\bigcap_{n \in G} A'_n \cap \bigcap_{n \in B} A'_n \neq \emptyset$ . It is easily seen that a Boolean independent sequence  $(A_n)$  of subsets of  $S$  has the property that  $(\chi_{A_n})$  is equivalent to the usual basis of  $l^1$  (see *Proposition 4*).

Because of the technical difficulties encountered in deducing *Theorem 1* from the above special case, we need a generalization of the notion of a convergent sequence of sets. It is also convenient to introduce the following terminology:

By a sequence we shall mean a set of objects indexed by some infinite subset  $M$  of the positive integers  $N$ ; we use the notation  $(f_n)_{n \in M}$ . We shall understand by "a subset of  $M$ " an infinite subset of  $M$ , unless the contrary is explicitly stated. Given  $L$  and  $M$  subsets of  $N$ , we say that  $L$  is almost contained in  $M$  if  $L \cap \sim M$  is a finite set. Given a sequence  $(f_n)_{n \in M}$  and subsets  $L$  and  $Q$  of  $M$  with  $L$  almost contained in  $Q$ , we call  $(f_n)_{n \in L}$  a subsequence of  $(f_n)_{n \in Q}$ . In the case in which  $(f_n)_{n \in M}$  is a sequence of real-valued functions defined on a set  $S$ , letting  $\{m_1, m_2, \dots\}$  be a strictly increasing enumeration of  $M$  and  $s \in S$ , we let

$$\overline{\lim}_M f_n(s) = \overline{\lim}_{j \rightarrow \infty} f_{m_j}(s) \text{ and } \lim_M f_n(s) = \lim_{j \rightarrow \infty} f_{m_j}(s).$$

(The point of our terminology, of course, is to avoid explicitly enumerating such sets  $M$  whenever it is feasible.)

**Definition:** Let  $S$  be a set,  $(A_n, B_n)_{n \in M}$  be a sequence of pairs of subsets of  $S$  with  $A_n \cap B_n = \emptyset$  for all  $n$ , and  $X$  a subset of  $S$ . We say that  $(A_n, B_n)_{n \in M}$  converges on the set  $X$  if every point  $x \in X$  either belongs to at most finitely many  $A_n$ 's, or to at most finitely many  $B_n$ 's, i.e., either  $\lim_{n \rightarrow \infty} \chi_{A_n}(x) = 0$  or  $\lim_{n \rightarrow \infty} \chi_{B_n}(x) = 0$ . (When  $X = S$ , the qualifier "on the set  $X$ " may be omitted.) We say that  $(A_n, B_n)_{n \in M}$  is independent if for every pair of disjoint finite nonempty subsets  $G$  and  $B$  of  $M$ ,

$$\bigcap_{n \in G} A_n \cap \bigcap_{n \in B} B_n \neq \emptyset. \quad [2]$$

We note that in the special case where  $B_n = S \sim A_n$  for all  $n$  and  $X = S$ ,  $(A_n, B_n)_{n \in M}$  converges on  $X$  if and only if  $(A_n)_{n \in M}$  converges. We also note that if  $(A_n, B_n)_{n \in M}$  converges on  $X$  and  $(A_n, B_n)_{n \in L}$  is a subsequence of  $(A_n, B_n)_{n \in M}$ , then  $(A_n, B_n)_{n \in L}$  converges on  $X$ ; if  $(A_n, B_n)_{n \in M}$  converges on each of the sets  $X_1, X_2, \dots$ , then  $(A_n, B_n)_{n \in M}$  converges on  $\bigcup_{i=1}^{\infty} X_i$ . Finally, it is an artifact of our definition

that every such sequence  $(A_n, B_n)_{n \in M}$  converges on the empty set. The special case of *Theorem 1* mentioned above, is an immediate consequence of the next result.

**THEOREM 2.** Let  $(A_n, B_n)_{n \in N}$  be a sequence of pairs of subsets of a set  $S$  with  $A_n \cap B_n = \emptyset$  for all  $n$ , and suppose that

$(A_n, B_n)_{n \in N}$  has no convergent subsequence. Then there is an infinite subset  $M$  of  $N$  so that  $(A_n, B_n)_{n \in M}$  is independent.

Theorem 2, in turn, follows very simply from the next crucial result.

LEMMA 3. Let  $l \geq 1$ ,  $(A_n, B_n)_{n \in N}$  a sequence of pairs of subsets of a set  $S$  with  $A_n \cap B_n = \phi$  for all  $n$ ,  $X_1, \dots, X_l$  disjoint subsets of  $S$ , and suppose that for each  $i$ ,  $1 \leq i \leq l$ ,  $(A_n, B_n)_{n \in N}$  has no subsequence convergent on  $X_i$ . Then there exists a  $j$  and an infinite subset  $M$  of  $N$  so that for each  $i$ ,  $1 \leq i \leq l$ ,  $(A_n, B_n)_{n \in M}$  has no subsequence convergent on  $X_i \cap A_j$  and also no subsequence convergent on  $X_i \cap B_j$ .

We introduce the following notation: Let  $n \in N$  and  $\epsilon = \pm 1$ ; define  $\epsilon A_n = A_n$  if  $\epsilon = +1$  and  $\epsilon A_n = B_n$  if  $\epsilon = -1$ .

We may deduce Theorem 2 from Lemma 3 by the following inductive process: Applying Lemma 3 for the case  $l = 1$ , choose  $n_1$  and  $M_1$  a subset of  $N$  so that  $(A_n, B_n)_{n \in M_1}$  has no subsequence convergent on either  $A_{n_1}$  or  $B_{n_1}$ . Suppose  $n_1 < n_2 < \dots < n_k$  and  $M_k$  have been chosen, so that on each of the  $2^k$  disjoint sets  $\bigcap_{j=1}^k \epsilon_j A_{n_j}$ ,  $(A_n, B_n)_{n \in M_k}$  has no convergent subsequence, where  $\epsilon = (\epsilon_1, \dots, \epsilon_k)$  ranges over all  $2^k$  choices of signs  $\epsilon_i = \pm 1$  all  $i$ . Now applying Lemma 3 for the case  $l = 2^k$ , choose  $n_{k+1} \in M_k$ ,  $n_{k+1} > n_k$ , and  $M_{k+1}$  a subset of  $M_k$  so that for each  $\epsilon = (\epsilon_1, \dots, \epsilon_k)$ ,  $(A_n, B_n)_{n \in M_{k+1}}$  has no subsequence convergent on  $\bigcap_{j=1}^k \epsilon_j A_{n_j} \cap A_{n_{k+1}}$  and also no subsequence convergent on  $\bigcap_{j=1}^k \epsilon_j A_{n_j} \cap B_{n_{k+1}}$ . This completes the definition of the  $n_j$ 's and  $M_j$ 's by induction; it now follows immediately that  $M = \{n_1, n_2, \dots\}$  satisfies the conclusion of Theorem 2. We note, incidentally, that the sequence  $(A_n, B_n)_{n \in M}$  has the property that it has no subsequence convergent on any nonempty member of the Boolean ring generated by  $\{A_n, B_n : n \in M\}$ .

We pass now to the proof of Lemma 3. The case  $l = 1$  is critical, and its proof is constructive in a certain sense. We shall exhibit an algorithm to produce the desired  $j$  and  $M$ ; this algorithm is then used to prove Lemma 3 by induction. We now suppose  $(A_n, B_n)_{n \in N}$  as in the Definition and let  $X$  be as subset of  $S$  so that  $(A_n, B_n)_{n \in N}$  has no subsequence convergent on  $X$ . We shall say that  $j$  and  $M$  work if  $(A_n, B_n)_{n \in M}$  has no subsequence convergent on either  $X \cap A_j$  or  $X \cap B_j$ . It is obvious that we can assume without loss of generality that  $S = X$ ; we do so.

THE BASIC ALGORITHM. Let  $n_1$  be an arbitrary element of  $N$ . If  $n_1$  and  $N$  do not work, let  $N_1$  be an arbitrary subset of  $N$  with the property that  $(A_n, B_n)_{n \in N_1}$  converges on  $A_{n_1}$  or  $B_{n_1}$ . Suppose  $k > 1$  and the subset  $N_{k-1}$  of  $N$  and the element  $n_{k-1}$  of  $N$  have been defined. Let  $n_k$  be an arbitrary member of  $N_{k-1}$  with  $n_k > n_{k-1}$ . If  $n_k$  and  $N_{k-1}$  do not work, let  $N_k$  be an arbitrary subset of  $N_{k-1}$  with the property that  $(A_n, B_n)_{n \in N_k}$  converges on  $A_{n_k}$  or  $B_{n_k}$ . We now assert that this process can be continued only a finite number of times. That is, as long as the  $n_j$ 's and  $N_j$ 's are selected in the above manner, there must exist a  $k \geq 1$  so that  $n_k$  works for  $N_{k-1}$  (where  $N_0 = N$ ).

Proof of this assertion: If not, we obtain  $n_k, N_k$ , and  $\epsilon_{n_k} = \pm 1$ , defined for all  $k \in N$ , with the properties that  $n_k > n_{k-1}$ ,  $n_k \in N_{k-1}$ ,  $N_k$  is a subset of  $N_{k-1}$ , and  $(A_n, B_n)_{n \in N_k}$  converges on  $\epsilon_{n_k} A_{n_k}$  (where we put  $n_0 = 0$ ).

Now let  $M = \{n_1, n_2, \dots\}$ . Then for every  $k$ ,  $(A_n, B_n)_{n \in M}$  is a subsequence of  $(A_n, B_n)_{n \in N_k}$ , hence  $(A_n, B_n)_{n \in M}$  con-

verges on  $\bigcup_{k=1}^{\infty} \epsilon_{n_k} A_{n_k}$ . Now we may choose an infinite subset  $M'$  of  $M$  so that  $\epsilon_m = 1$  for all  $m \in M'$ , or  $\epsilon_m = -1$  for all  $m \in M'$ . Suppose the first possibility. Since  $(A_n, B_n)_{n \in M'}$  is a subsequence of  $(A_n, B_n)_{n \in M}$  and  $\bigcup_{k=1}^{\infty} \epsilon_{n_k} A_{n_k} \supset \bigcup_{n \in M'} \epsilon_n A_n = \bigcup_{n \in M'} A_n$ ,  $(A_n, B_n)_{n \in M'}$  converges on  $\bigcup_{n \in M'} A_n$ . Since  $(A_n, B_n)_{n \in M'}$  does not converge, there must exist an  $x$  so that  $\{n \in M' : x \in A_n\}$  is infinite and also  $\{n \in M' : x \in B_n\}$  is infinite. But then  $x \in \bigcup_{n \in M'} A_n$ , and hence  $(A_n, B_n)_{n \in M'}$  does not converge on  $\bigcup_{n \in M'} \epsilon_n A_n$ , a contradiction. The proof for the case of the second possibility is the same.

The proof of the assertion of the Basic Algorithm, and hence of the case  $l = 1$  of Lemma 3, is now complete. Suppose Lemma 3 proved for  $l = r$ , and let the  $X_i$ 's and  $(A_n, B_n)_{n \in N}$  satisfy its hypotheses for the case  $l = r + 1$ . Again, we shall say that  $j$  and  $M$  work if  $(A_n, B_n)_{n \in M}$  has no subsequence convergent on either  $X_{r+1} \cap A_j$  or  $X_{r+1} \cap B_j$ . We shall also say that  $j$  and  $M$   $r$ -work if for every  $1 \leq i \leq r$  and  $\epsilon = \pm 1$ ,  $(A_n, B_n)_{n \in M}$  has no subsequence convergent on  $X_i \cap \epsilon A_j$ . By the induction hypothesis, we may choose  $n_1$  and  $N'_1$  a subset of  $N$  so that  $n_1$  and  $N'_1$   $r$ -work. If  $n_1$  and  $N'_1$  do not work, choose  $N_1$  a subset of  $N'_1$  so that  $(A_n, B_n)_{n \in N_1}$  converges on  $A_{n_1} \cap X_{r+1}$  or on  $B_{n_1} \cap X_{r+1}$ . Suppose  $k > 1$ , and the subset  $N_{k-1}$  of  $N$  and the element  $n_{k-1}$  of  $N$  have been defined. Since  $(A_n, B_n)_{n \in N_{k-1}}$  is a subsequence of  $(A_n, B_n)_{n \in N}$ , we may apply the induction hypothesis to choose an  $n_k \in N_{k-1}$  with  $n_k > n_{k-1}$  and a subset  $N'_k$  of  $N_{k-1}$  so that  $n_k$  and  $(A_n, B_n)_{n \in N'_k}$   $r$ -work. Again, if  $n_k$  and  $(A_n, B_n)_{n \in N'_k}$  do not work, choose  $N_k$  a subset of  $N'_k$  so that  $(A_n, B_n)_{n \in N_k}$  converges on  $A_{n_k} \cap X_{r+1}$  or on  $B_{n_k} \cap X_{r+1}$ . Now this process cannot be continued indefinitely, since the  $n_k$ 's and  $N_k$ 's thus constructed satisfy the criteria of the Basic Algorithm and  $(A_n, B_n)_{n \in N}$  has no subsequence convergent on  $X_{r+1}$ . Thus, there must exist a  $k \geq 1$  so that  $n_k$  and  $N'_k$  work. By construction  $n_k$  and  $N'_k$   $r$ -work, hence by definition,  $n_k$  and  $N'_k$  satisfy the conclusion of Lemma 3.

This completes the proof of Lemma 3 and hence of Theorem 2. To apply Theorem 2 to the proof of our Main Theorem, we need the following simple sufficient condition for a sequence of functions to be equivalent to the usual  $l^1$ -basis.

PROPOSITION 4: Let  $(f_n)_{n \in M}$  be a uniformly bounded sequence of real-valued functions defined on a set  $S$  and  $\delta$  and  $r$  real numbers with  $\delta > 0$ . Assume, putting  $A_n = \{x : f_n(x) > \delta + r\}$  and  $B_n = \{x : f_n(x) < r\}$  for all  $n \in M$ , that  $(A_n, B_n)_{n \in M}$  is independent. Then  $(f_n)_{n \in M}$  is equivalent, in the supremum norm, to the usual  $l^1$ -basis.

Proof: We shall prove that the " $\delta$ " of Eq. [1] may be chosen to be  $\delta/2$ . By multiplying all the  $f_n$ 's by  $-1$  if necessary, we may assume that  $\delta + r > 0$ . Let  $(c_i)_{i \in M}$  be a sequence of scalars with only finitely many  $c_i$ 's non-zero and  $\sum |c_i| = 1$ . It suffices to show that there is an  $s$  in  $S$  with

$$|\sum c_i f_i(s)| \geq \delta/2. \quad [3]$$

Let  $G = \{i \in M : c_i > 0\}$  and  $B = \{i \in M : c_i < 0\}$ . Since Eq. [2] holds, we may choose  $x$  and  $y$  such that  $x \in \bigcap_{i \in G} A_i \cap \bigcap_{i \in B} B_i$  and  $y \in \bigcap_{i \in B} A_i \cap \bigcap_{i \in G} B_i$ . If we suppose first that  $r \geq 0$  and set  $B' = \{i \in B : f_i(x) > 0\}$ , then

$$\sum_{i \in B} c_i f_i(x) \geq \sum_{i \in B'} c_i f_i(x) > -r \sum_{i \in B'} |c_i| \geq \sum_{i \in B} |c_i| (-r). \quad [4]$$

Similarly,

$$-\sum_{i \in G} c_i f_i(y) \geq \sum_{i \in G} |c_i|(-r). \quad [5]$$

By Eqs. [4], [5], and the definitions of  $x$  and  $y$ , we thus have

$$\sum c_i f_i(x) \geq \sum_{i \in G} |c_i|(\delta + r) + \sum_{i \in B} |c_i|(-r) \quad [6]$$

and

$$-\sum c_i f_i(y) \geq \sum_{i \in B} |c_i|(\delta + r) + \sum_{i \in G} |c_i|(-r). \quad [7]$$

It is easily seen that Eqs. [6] and [7] also hold if  $r < 0$ . Since the sum of the right-hand sides of Eqs. [6] and [7] equals  $\delta$ , the maximum of the left-hand sides must be at least as large as  $\delta/2$ . Here we tacitly assumed  $G \neq \emptyset$  and  $B \neq \emptyset$ ; however, the argument is still valid if we simply replace an intersection over the empty set of indices by  $S$ , and a sum over the empty set of indices by 0. Thus, Eq. [3] is established for  $s = x$  or  $s = y$ , so Proposition 4 is proved.

We now assume that  $S$  and  $(f_n)_{n \in M}$  satisfy the hypotheses of Theorem 1 and fail the first alternative; i.e.  $(f_n)_{n \in N}$  has no subsequence point-wise convergent on  $S$ . To complete the proof of Theorem 1, it suffices to construct  $\delta > 0$ , a real number  $r$ , and a subset  $M$  of  $N$  so that the hypotheses of the preceding proposition are satisfied. The next two lemmas allow us to find  $\delta$  and  $r$ ; their demonstrations involve standard arguments.

LEMMA 5. For each subset  $M$  of  $N$ , let

$$\delta(M) = \sup_{x \in S} (\lim_M f_m(x) - \lim_M f_m(x)).$$

Then there exists a subset  $Q$  of  $N$  so that for all subsets  $L$  or  $Q$ ,  $\delta(L) = \delta(Q)$ .

Remark: Our standing assumptions imply that  $\delta(M) > 0$  for all subsets  $M$  of  $N$ .

Proof of Lemma 5: For any subsets  $L$  and  $M$  of  $N$  with  $L$  almost contained in  $M$ , we have that  $\delta(L) \leq \delta(M)$ . Were the conclusion of Lemma 5 false, we could choose by transfinite induction, a transfinite family  $\{N_\alpha: \alpha < \omega_1\}$  of subsets of  $N$ , indexed by the set of ordinals  $\alpha$  less than the first uncountable ordinal  $\omega_1$ , with the property that for all  $\alpha < \beta < \omega_1$ ,  $N_\beta$  is almost contained in  $N_\alpha$  and  $\delta(N_\beta) < \delta(N_\alpha)$ . This is impossible, for as is well known, there does not exist a transfinite strictly decreasing sequence of (positive) real numbers. (To reach a contradiction, simply put  $\delta = \inf \{\delta(N_\alpha): \alpha < \omega_1\}$ , then choose  $(\alpha_n)_{n \in N}$  a sequence of ordinals with  $\delta = \lim \delta(N_{\alpha_n})$ ; for  $\beta > \sup_N \alpha_n$ , we have  $\delta(N_\beta) < \delta$ ).

Now choose  $Q$  a subset of  $N$ , satisfying the conclusion of Lemma 5, and put  $\delta = \delta(Q)/2$ ; by the above remark,  $\delta > 0$ .

LEMMA 6. There exists a subset  $M'$  of  $Q$  and a rational number  $r$  so that for every subset  $L$  of  $M'$ , there is an  $x \in S$  satisfying

$$\lim_L f_i(x) > \delta + r \text{ and } \lim_L f_i(x) < r.$$

Theorem 1 and, hence, our main result follow immediately from Theorem 2, Proposition 4, and Lemma 6. Indeed, let  $M'$  satisfy the conclusion of Lemma 6, and for each  $n \in M'$ , let  $A_n = \{x \in S: f_n(x) > \delta + r\}$  and  $B_n = \{x \in S: f_n(x) < r\}$ . The conclusion of Lemma 6 yields that  $(A_n, B_n)_{n \in M'}$  has no convergent subsequence. By Theorem 2, we may select a subset  $M$  of  $M'$  so that  $(A_n, B_n)_{n \in M}$  is independent. Then

$(f_n)_{n \in M}$  satisfies the hypotheses of Proposition 4, hence is equivalent in the supremum norm, to the usual  $l^1$ -basis.

Proof of Lemma 6: Suppose not. Let  $r_1, r_2, \dots$  be an enumeration of the rational numbers. Choose  $L_1$  a subset of  $Q$  so that

$$\text{for all } x \in S, \lim_L f_i(x) \leq \delta + r \text{ or } \lim_L f_i(x) \geq r \quad [8]$$

holds for  $L = L_1$  and  $r = r_1$ . Having chosen the subset  $L_k$  of  $Q$ , choose  $L_{k+1} \subset L_k$  so that Eq. [8] holds for  $L = L_{k+1}$  and  $r = r_{k+1}$ . This defines  $L_1 \supset L_2 \supset \dots \supset L_k \dots$  by induction. Now by the standard diagonal procedure, choose an infinite set  $L$  with  $L$  almost contained in  $L_k$  for all  $k \in N$ . It then follows that Eq. [8] holds for all rational numbers  $r$ . Since  $L$  is in turn almost contained in  $Q$  and  $Q$  satisfies the conclusion of Lemma 5,  $\delta(L) = \delta(Q) = 2\delta$ . Let  $\epsilon = \delta/2$ . By the definition of  $\delta(L)$ , we may choose an  $x \in S$  so that

$$\lim_L f_i(x) - \lim_L f_i(x) > \delta(L) - \epsilon. \quad [9]$$

Now let  $a = \lim_L f_i(x)$  and  $b = \lim_L f_i(x)$ ; Eq. [9] may then be expressed in the form

$$a > 2\delta - \epsilon + b > b.$$

Choose a rational number  $r$  so that  $r > b$  and  $r - b + \delta < 2\delta - \epsilon = (3/2)\delta$ . Thus

$$b < r < r + \delta = (r - b) + \delta + b < 2\delta - \epsilon + b < a.$$

Since we thus have  $a > \delta + r$  and  $b < r$ , Eq. [8] is contradicted.

Q.E.D.

We do not know if the Main Theorem holds for complex Banach spaces\*. However let  $B$  be a complex Banach space, and  $(e_n)_{n \in N}$  a sequence in  $B$  equivalent to the usual  $l^1$ -basis over the real scalars. If there is no subset  $M$  of  $N$  with  $(e_n)_{n \in M}$  equivalent to the usual  $l^1$ -basis over the complex scalars, we may choose finite disjoint subsets  $B_1, B_2, \dots$  of  $N$  and elements  $y_1, y_2, \dots$  and  $z_1, z_2, \dots$  in  $B$  so that  $\lim_{j \rightarrow \infty} \|y_j - z_j\| = 0$  so that for all  $j$ ,  $\|y_j\| = \|z_j\| = 1$  and  $y_j$  and  $iz_j$  are in the linear span of  $\{e_n: n \in B_j\}$ . Now for all real scalars  $a$  and  $b$  and all  $j$ , we have that

$$\|aiz_j + by_j\| \geq \|(ai + b)z_j\| - \|by_j - bz_j\| \\ = (a^2 + b^2)^{1/2} - |b| \|y_j - z_j\|.$$

It then follows that there is a  $k$  so that both sequences  $(y_k, iz_k, y_{k+1}, iz_{k+1}, \dots)$  and  $(z_k, iz_k, z_{k+1}, iz_{k+1}, \dots)$  are equivalent to the usual  $l^1$ -basis over the real scalars. Consequently  $(z_k, z_{k+1}, \dots)$  is equivalent to the usual  $l^1$ -basis over the complex scalars. Hence our Main Theorem implies that the result stated in the abstract holds for complex Banach spaces as well.

This research was partially supported by NSF-GP 30798X1.

Addendum. E. Odell and I have shown that a separable Banach space  $B$  contains a subspace isomorphic to  $l^1$  provided there exists an element in  $B^{**}$  that is not a limit in the  $B^*$  topology of a sequence in  $B$ . Consequently,  $B$  contains an isomorph of  $l^1$  if (and only if) the cardinality of  $B^{**}$  is greater than that of the continuum. The proof uses Proposition 4 and arguments similar to those of Lemmas 5 and 6, but does not make use of Theorem 2 or Lemma 3. This result and related ones will appear elsewhere.

\* Note Added in Proof. This has been resolved in the affirmative by L. Dor.